

## Expression of CR1 (CD35) mRNA in podocytes from adult and fetal human kidneys

MARIE-DOMINIQUE APPAY, MICHEL D. KAZATCHKINE, MATTHIEU LEVI-STRAUSS,  
NICOLE HINGLAIS, and JEAN BARIETY

*INSERM U28, Hôpital Broussais, and INSERM U276, Institut Pasteur, Paris, France*

**Expression of CR1 (CD35) mRNA in podocytes from adult and fetal human kidneys.** The presence of CR1 mRNA in podocytes was investigated using a  $^{35}\text{S}$ -labeled CR1 cDNA probe and in situ hybridization in sections from fetal and adult human kidneys. CR1 mRNA was only detected in immature podocytes at early stages of glomerular differentiation in the fetal kidney. In contrast, CR1 antigen was abundantly expressed on immature and mature podocytes in fetal kidneys and adult glomeruli. Thus, the expression of the CR1 gene in podocytes is tightly regulated. It is possible that the prolonged life span of adult podocytes is associated with a slow turnover of CR1 and low or intermittent accumulation of CR1 mRNA transcripts.

The human complement receptor type 1 (CR1, CD35) is a polymorphic single chain membrane glycoprotein expressed on human erythrocytes, neutrophils, monocytes/macrophages, B lymphocytes, a subset of CD4<sup>+</sup> T lymphocytes, follicular dendritic cells, Kupffer cells and human adult and fetal podocytes [1–12]. By immunoelectron microscopy, CR1 antigen has been observed on the plasma membrane, in coated pits [13] and in the Golgi apparatus [6] of adult podocytes. The number of CR1 molecules expressed in adult podocytes has been estimated to be  $2 \cdot 10^5$  per cell [14]. Using in situ hybridization, we have now examined the presence of specific CR1 mRNA in podocytes from adult and fetal human kidneys. CR1 mRNA-containing podocytes were present in immature glomeruli from fetal kidney, whereas no CR1 mRNA-containing podocytes were detected in adult kidney sections. This is the first example, in the human kidney, of the presence of a specific mRNA in cells during ontogenesis which becomes undetectable in normal mature glomeruli, although the antigenic expression of the gene product persists.

### Methods

#### *Tissues*

Normal adult kidney tissue specimens were obtained from normal parts from two kidneys immediately after surgical

nephrectomy for renal cell carcinoma. Human fetal kidneys were obtained from 20 and 26 week pregnancies according to the guidelines of the French National Committee of Ethics. Tonsil tissues were obtained following elective surgery. Tissue specimens were immediately quick-frozen and stored in liquid nitrogen until sectioned.





The 0.75 kb CR1-1 cDNA probe from human tonsil [15] was from D.T. Fearon (Johns Hopkins Medical School, Baltimore, Maryland, USA). The insert was excised from the plasmid using Eco R1 (Appligene, Illkirch, France) and isolated by electrophoresis prior to labeling. For in situ hybridization, the CR1 cDNA was labeled with  $^{35}\text{S}$  dCTP (Amersham, Les Ulis, France) using random oligonucleotides as primers to a specific activity of 0.9 to  $2 \cdot 10^9$  cpm/ $\mu\text{g}$ . The labeled probe was ethanol-precipitated and resuspended at 10  $\mu\text{g}/\text{ml}$  in 10 mM Tris-HCL, 1 mM EDTA, pH 7.5. Controls were a 0.9 kb fragment of the recombinant plasmid, and a renin cDNA (from Dr. P. Corvol, INSERM U36, Paris, France) labeled to the same specific activity.

#### *In situ hybridization*

In situ hybridization was performed as previously reported [16] with minor modifications. Briefly, frozen tissue sections fixed in 3.7% paraformaldehyde in phosphate buffer saline (PBS) were sequentially digested with 0.2 N HCl and 10  $\mu\text{g}/\text{ml}$  proteinase K for 30 minutes at 37°C. After dehydration and air-drying,  $^{35}\text{S}$ -labeled CR1 cDNA probe ( $10^6$  cpm/slide) in 15  $\mu\text{l}$  of hybridization mixture was applied to sections. The hybridizing mixture contained: 50% Dowex AG deionized formamide, 10% dextran, 10 mM Tris pH 7.4, 1 mM EDTA, NaCl 0.6 N, 1 $\times$  Denhardt solution, 10 mM DTT, yeast tRNA (10 mg/ml), sheared salmon sperm cDNA (400  $\mu\text{g}/\text{ml}$ ), and herring sperm DNA (400  $\mu\text{g}/\text{ml}$ ). The sections were covered with siliconized glass coverslips and sealed. The slides were incubated, for 18 hours at 37°C, de-coverslipped and washed in a solution containing 50% formamide and 50% 8 $\times$ SSC for 15 minutes at 20°C, twice in 2 $\times$ SSC for 30 minutes at 20°C and in 2 $\times$ SSC for 24 hours at 20°C under gentle agitation. Slides were dehydrated in graded alcohols before being dipped into a 50% NTB2 emulsion (Kodak) in water. Autoradiography was for two weeks (human tonsils) and eight weeks (kidneys) in the dark at 4°C. Autoradiographies were developed in Kodak D19 (1.5 to 3 minutes,

Received for publication August 7, 1989  
and in revised form February 21, 1990  
Accepted for publication February 22, 1990

© 1990 by the International Society of Nephrology



**Fig. 1.** *In situ* hybridization with the  $^{35}\text{S}$  CR1-1 DNA probe of a human fetal kidney section (20 gestation weeks). Hybridizing cells are present at the periphery of an immature glomerulus at the open glomerulus maturation stage. ( $\times 500$ )

**Fig. 2.** Immunoperoxidase detection of CR1 antigen in a semithin section of a human fetal kidney using  $\text{Fab}'_2$  rabbit anti-CR1 antibodies and peroxidase-labeled  $\text{Fab}'_2$  sheep anti-rabbit antibodies. The podocyte membrane is stained by the reaction product of peroxidase labeled second antibody ( $\times 500$ ). The glomerulus is at the same stage of differentiation as that shown in Fig. 1. Similar results were obtained using the monoclonal anti-CR1 antibody J3D3 [20] (Data not shown).

**Fig. 3.** *In situ* hybridization with the CR1 cDNA CR1-1 probe of an adult human kidney section. No mRNA-containing cell is detectable in the glomerulus. ( $\times 250$ ). Insert: detailed view of the same glomerulus ( $\times 500$ ).

**Fig. 4.** Indirect immunofluorescence detection of CR1 antigen in a cryostat section of a human adult kidney using  $\text{Fab}'_2$  rabbit anti-CR1 antibodies and FITC-labeled sheep anti-rabbit antibodies. The CR1 antigen is detected in all the podocytes of the glomerulus ( $\times 250$ ).

20°C) and fixed in Kodak rapid fix for five minutes (20°C). As a final step, the slides were stained with hematoxylin and eosin.

Negative control experiments included 1) pretreatment with RNase A (Sigma) at a concentration of 40  $\mu\text{g}/\text{ml}$  in  $2\times\text{SSC}$  for 30 minutes at 37°C prior hybridization, 2) hybridization of some slides with  $^{35}\text{S}$ -labeled recombinant plasmid, 3) hybridization of some slides with  $^{35}\text{S}$ -labeled renin cDNA.

A positive control experiment was performed by incubating frozen sections of human tonsils with CR1 cDNA probe [15].

#### *Quantitation of in situ hybridization signal by image analysis*

Image analysis was performed on CR1 probe-hybridized sections from two fetal kidneys and two adult kidneys, and on one RNase A-treated fetal kidney section. We used a computer

vision image processor (NS 1500, Nachet-Vision, France), designed according to the "mathematical morphology" theory [17]. Silver grains segmentation was easy at the magnification  $\times 100$ . Silver grains were isolated by "top hat" transformation, to select them according to their size and contrast [18, 19]. Because of the silver grain image overlapping, their surface rather than their absolute number was calculated. For silver grains localization, the operator displaced a circular mask upon immature glomerular structures, mature glomerular structures and tubular structures. The results were expressed per area unit (100  $\mu\text{m}^2$ ). Five measurements were performed on twenty immature glomeruli, on ten mature glomeruli from fetal kidneys and five measurements were performed on ten glomeruli of adult kidney. A Newman-Keuls test was performed in a one

way analysis of variance on measurements. Results were presented as histograms.

### Immunohistochemistry

Alternate cryostat sections of specimens were air dried and acetone fixed for 10 minutes at room temperature for the detection of CR1 antigen by indirect immunofluorescence, using mouse monoclonal or polyclonal rabbit antibodies against CR1 [6, 20]. The second antibody was either goat anti-mouse fluoresceinated antibody or sheep anti-rabbit fluoresceinated antibody (Biosys, Compiègne, France). Some fetal kidney specimens were stained with Fab'<sub>2</sub> anti-CR1 polyclonal antibodies [6] followed by peroxidase-labeled Fab'<sub>2</sub> sheep anti-rabbit IgG, and processed for immunoelectron microscopy as previously described [11].

## Results

### Fetal and adult human kidneys

CR1 mRNA-containing cells were detected in immature glomeruli from fetal kidneys from the stage of open glomerulus to that of plurilobulated glomerulus. No hybridizing cells were detected in fully lobulated glomeruli, that is, in the most mature stage of glomeruli. In glomeruli at the open glomerulus stage, the mRNA-containing cells appeared to be ring-like, and were distributed at the periphery of the glomeruli in the area where immature podocytes were present [11] (Fig. 1). The distribution of hybridizing cells was closely related to that of CR1 antigen at that stage (Fig. 2). In glomeruli at the plurilobulated glomerulus stage, the number of mRNA-containing podocytes was lower than that of CR1-immunoreactive podocytes (data not shown). In mature fetal glomeruli and in adult kidney sections, no mRNA-containing cells were detected, although CR1 antigen was abundantly expressed on podocytes (Figs. 3 and 4).

Image analysis confirmed in a quantitative fashion that CR1 mRNA was significantly more abundant in immature glomeruli than in mature fetal glomeruli and that no CR1 mRNA was detected in mature glomeruli from fetal kidneys nor in glomeruli from adult kidneys (Fig. 5). No CR1 mRNA was present in tubular cells in fetal and adult kidneys.

### Controls


No CR1 cDNA hybridization was observed in fetal kidney sections that had been pretreated with RNase.

No hybridizing cells were detected in adult and fetal tissues using the control plasmid probe or the renin cDNA probe.

Human tonsils served as a positive control for CR1 cDNA hybridization. CR1 cDNA-hybridizing cells in human tonsils were distributed in the mantle zone and in the center of the secondary follicles (Fig. 6). The distribution of CR1 cDNA-hybridizing cells correlated with that of CR1 antigen-expressing cells.

## Discussion

C3b CR1 is abundantly expressed on the plasma membrane of adult human podocytes [6, 14]. CR1 antigen is an early marker of podocyte differentiation in human fetal kidneys [11]. In adult podocytes, the CR1 molecule is functional in that it can bind C3b, which is attached to surfaces, and decay-dissociate pre-formed alternative pathway C3 convertase complexes [7-14].



**Fig. 5.** Quantitative image analysis of silver grains (grains surface per surface unit) in CR1 cDNA CR1-1-hybridized kidney sections. The histograms depict the mean calculated surface of grains per 100  $\mu\text{m}^2$  in immature glomeruli (Im. Glo.), mature glomeruli (M. Glo.), tubules (Tub.) and in RNase-treated (control) immature glomeruli (RNase Im. Glo.) from two fetal kidneys (A) and in glomeruli (Glo.) and tubules (Tub.) from two adult kidneys (B), with 95% confidence interval.

The physiological role of CR1 on podocytes remains, however, unclear. The present study investigated the synthesis of the CR1 protein in fetal and adult human kidney by in situ hybridization.

In fetal kidneys the CR1 mRNA-containing cells with the CR1-1 cDNA probe was detected in early stages of glomerular maturation. Immunofluorescence and in situ hybridization studies indicated that CR1 is both expressed and actively synthesized by immature podocytes. CR1 mRNA expression decreased with the maturation of glomeruli. Thus, on the same cryostat section of fetal kidney, CR1 mRNA was present in immature glomeruli at the open and plurilobulated stages, and was not detected in mature (fully lobulated) glomeruli.

In adult kidney, the CR1 antigen was abundantly present on the plasma membrane of podocytes [6, 14]. However, no CR1 mRNA was detected in the cells. The latter observation is consistent with the lack of detectable CR1 mRNA in mature glomeruli in fetal kidneys. The absence of CR1 mRNA in podocytes of adult kidneys was unexpected in the light of the previous finding of CR1 antigen in the endoplasmic reticulum






Fig. 6. In situ hybridization with the  $^{35}\text{S}$  CR1-1 DNA probe of a human tonsil. Hybridizing cells are present in the germinal center (arrow) and at the periphery of the germinal center (double arrow) ( $\times 280$ ).

and endocytic vesicles of adult podocytes [6, 13]. It is possible that CR1 mRNA is present in adult podocytes in amounts that are below the detection threshold of in situ hybridization techniques. A long exposure of two months was required in order to detect CR1 mRNA in fetal kidney sections, suggesting that detectable CR1 mRNA levels in the kidney are low, in contrast with human tonsils in which a ten day exposure time was sufficient for CR1 mRNA detection. It is also possible that the turnover of the CR1 mRNA in adult glomeruli occurs intermittently, or alternatively, that it is very slow and that only few CR1 mRNA transcripts are sufficient to ensure the podocyte requirement for new CR1 glycoprotein molecules. Consistent with the latter hypothesis is the observation of a prolonged life span of podocytes in normal rats [21]. Newly synthesized CR1 glycoprotein would accumulate in the Golgi apparatus and may be detected by immunoelectron microscopy. The mere detection of CR1 antigen on the plasma membrane in the Golgi apparatus and in endocytic vesicles is not indicative of ongoing synthesis of the protein, since it could also reflect the intracellular traffic of the protein between the plasma membrane and the Golgi apparatus [22]. In recent studies, it was observed that specific mRNA for human elastase was detected in cells at specific differentiation stages in the myelocytic lineage. mRNA transcripts were absent in mature cells of the neutrophil myeloid series although the cells contained the protein [23].

Results of the present investigation indicate that the expression of the CR1 gene in podocyte is tightly controlled and differently regulated at various stages of cell differentiation.

We are currently examining whether CR1 mRNA may be expressed in glomeruli of diseased kidneys.

#### Acknowledgments

Renin cDNA was a gift from Dr. P. Corvol, INSERM U36, Paris, France. The CR1-1 cDNA probe from human tonsil used in this study

was a gift from D.T. Fearon, Johns Hopkins Medical School, Baltimore, Maryland, USA. We are grateful to Dr. Brossard from the Centre d'Hémodiologie Périnatale, Paris, for providing human fetal kidney samples used in this study. The authors thank Mr. M. Paing for photography.

Reprint requests to Marie-Dominique Appay, INSERM U28, Hôpital Broussais, rue Didot, Paris Cedex 14, France.

#### References

1. FEARON DT: Identification of the membrane glycoprotein that is the C3b receptor of the human erythrocyte, PMN leukocyte, B-lymphocyte and monocyte. *J Exp Med* 152:20-30, 1980
2. WILSON JG, TEDDER TF, FEARON DT: Characterization of human T lymphocytes that express the C3b receptor. *J Immunol* 113: 684-689, 1983
3. COHEN JHM, CAUDWELL V, LEVI-STRAUSS M, BOURGEOIS P, KAZATCHKINE MD: Genetic analysis of CR1 expression on erythrocytes of patients with systemic lupus erythematosus. *Arth Rheum* 32:393-397, 1989
4. REYNES M, AUBERT JP, COHEN JHM, AUDOUIN J, TRICOTTET J, DIEBOLD J, KAZATCHKINE MD: Human follicular dendritic cells express CR1, CR2 and CR3 complement receptor antigens. *J Immunol* 135:2687-2694, 1985
5. HINGLAIS N, KAZATCHKINE MD, MANDET C, APPAY MD, BARIETY J: Human liver Küpfers cells express CR1, CR3 and CR4 complement receptor antigens. An immunohistochemical study. *Lab Invest* 61:509-514, 1990
6. KAZATCHKINE MD, FEARON DT, APPAY MD, MANDET C, BARIETY J: Immunohistochemical study of the human glomerular C3b receptor in normal kidney and in seventy-five cases of renal diseases. *J Clin Invest* 69:900-912, 1982
7. SHIN ML, GELFAND MC, NAGLE RB, CARLO JR, GREEN I, FRANK MM: Localization of receptors for activated complement on visceral epithelial cells of the human renal glomerulus. *J Immunol* 118:869-873, 1977
8. GELFAND MC, FRANCK MM, GREEN I: A receptor for the third component of complement in the human renal glomerulus. *J Exp Med* 145:1029-1040, 1975
9. MATRE R, NILSEN R, THUNOLD S, TONDER O: The localization of glomerular C3b receptor by immunoelectron microscopy. *Scand J Immunol* 16:351-354, 1982
10. EMANCIPATOR SN, IIDA K, NUSSENZWEIG V, GALLO GR: Monoclonal antibodies to human complement receptor (CR1) detect defects in glomerular diseases. *Clin Immunol Immunopathol* 27: 170-176, 1983
11. APPAY MD, MOUNIER F, GUBLER MC, ROUCHON M, BEZIAU A, KAZATCHKINE MD: Ontogenesis of the glomerular C3b receptor (CR1) in fetal human kidney. *Clin Immunol* 37:103-113, 1985
12. APPAY MD, KAZATCHKINE MD, MOUNIER F, BARIETY J: Presence of the C3b complement receptor (CR1) antigen in the glomerular basement membrane of human fetal kidneys. *Nephrol Dial Transplant* 2:162-165, 1988
13. WEISS L, FISCHER E, HAEFFNER-CAVAILLON N, JOUVIN MH, APPAY MD, BARIETY J, KAZATCHKINE MD: The human C3b receptor (CR1) in *Advances in Nephrology* edited by GRUNFELD, MAXWELL, BACH, FUNCK-BRENTANO, Chicago, London, Boca-Raton, Year Book Medical Publishers, 1989, p. 260
14. FISCHER E, APPAY MD, COOK J, KAZATCHKINE MD: Characterization of the human glomerular C3 receptor as the C3b/C4b complement type one (CR1) receptor. *J Immunol* 136:1373-1377, 1986
15. WONG WW, KLICKSTEIN LB, SMITH JH, WEIS JH, FEARON DT: Identification of a partial cDNA clone for the human receptor for complement fragments C3b/C4b. *Proc Natl Acad Sci USA* 87: 7711-7715, 1985
16. FOURNIE JG, TARDIEU M, LEBON P, ROBAIN O, PONSOT G, ROZENBLATT S, BOUTELLE M: Detection of measles virus RNA in lymphocytes from peripheral-blood and brain perivascular infiltrates of patients with subacute sclerosing panencephalitis. *N Engl J Med* 313:910-915, 1985

17. SERRA J: *Image Analysis and Mathematical Morphology*. London, Academic Press, 1982
18. MEYER F, SERRA J: Méthodes morphologiques de transformation des images biologiques. *13e Congrès des Anatomo-Pathologistes de Langue Française*, Paris, 1983, p. 2-19
19. SALZMANN JL, BERNIAU D, POLIARD A, BOUSSARD C, FELDMANN G: An automatic method for counting silver grains in autohistoradiography: An application to in situ hybridization on rat liver cells. *Acta Stereol* 5:273-278, 1986
20. COOK J, FISCHER E, BOUCHEIX C, MIRSAHI M, JOUVIN MH, WEISS L, JACK RM, KAZATCHKINE M: Mouse monoclonal antibodies to the human C3b receptor. *Mol Immunol* 22:531-539, 1985
21. PABST R, STERZEL RB: Cell renewal of glomerular cell types in normal rats. An autoradiographic analysis. *Kidney Int* 24:626-631, 1983
22. BRANDS RM, SNIDER D, HINO Y, PARK SS, GELBOIN HV, ROTHMAN JE: Retention of membrane proteins by the endoplasmic reticulum. *J Cell Biol* 101:1724-1732, 1985
23. FOURET P, DU BOIS RM, BERNAUDIN JF, TAKAHASHI H, FERRANS VJ, CRYSTAL RG: Expression of the neutrophil elastase gene during human bone marrow cell differentiation. *J Exp Med* 169:833-845, 1989